

On multivariate projection operators

L. Szili^a, P. Vértesi^{b,*}

^a *Department of Numerical Analysis, Loránd Eötvös University, Budapest, Pázmány P. sétány 1/C, H-1117, Hungary*

^b *Rényi Mathematical Institute of the Hungarian Academy of Sciences, Budapest, Reáltanoda u. 13–15, H-1053, Hungary*

Received 10 April 2008; accepted 24 November 2008

Available online 3 December 2008

Communicated by C.K. Chui and H.N. Mhaskar

Dedicated to the memory of G.G. Lorentz

Abstract

This paper deals with multivariate Fourier series considering triangular type partial sums. Among others we give the exact order of the corresponding operator norm. Moreover, a generalization of the so-called Faber–Marcinkiewicz–Berman theorem has been proved.

© 2008 Elsevier Inc. All rights reserved.

Keywords: Rectangular partial sums; Triangular partial sums; Operator norm of the multivariate Fourier series; Projection operators

1. Introduction

1.1

Multivariate Fourier series has been the object of an intensive study. We may refer the classical works Zygmund [1, Ch. XVII] and Stein, Weiss [2, Ch. VII]. This paper is another contribution to this subject. To step further we need some notations.

Let \mathbb{R}^d (direct product) be the Euclidean d -dimensional space ($d \geq 1$, fixed) and let $\mathbb{T}^d = \mathbb{R}^d \pmod{2\pi\mathbb{Z}^d}$ denote the d -dimensional torus, where $\mathbb{Z} = \{0, \pm 1, \pm 2, \dots\}$.

Further, let $C(\mathbb{T}^d)$ denote the space of (complex valued) continuous functions on \mathbb{T}^d . By definition they are 2π -periodic in each variable.

* Corresponding author.

E-mail addresses: szili@ludens.elte.hu (L. Szili), vertesi@renyi.hu, veter@renyi.hu (P. Vértesi).

For $g \in C(\mathbb{T}^d)$ we define its Fourier series by

$$g(\boldsymbol{\vartheta}) \sim \sum_{\mathbf{k}} \hat{g}(\mathbf{k}) e^{i\mathbf{k} \cdot \boldsymbol{\vartheta}}, \quad \hat{g}(\mathbf{k}) = \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} g(\mathbf{t}) e^{-i\mathbf{k} \cdot \mathbf{t}} d\mathbf{t}, \quad (1.1)$$

where in the above vector notation $\boldsymbol{\vartheta} = (\vartheta_1, \vartheta_2, \dots, \vartheta_d) \in \mathbb{T}^d$, $\mathbf{k} = (k_1, k_1, \dots, k_d) \in \mathbb{Z}^d$ and $\mathbf{k} \cdot \boldsymbol{\vartheta} = \sum_{l=1}^d k_l \vartheta_l$ (scalar product).

The *rectangular* n th partial sum of the Fourier series is defined by

$$S_{nd}^{[r]}(g, \boldsymbol{\vartheta}) := \sum_{|\mathbf{k}|_\infty \leq n} \hat{g}(\mathbf{k}) e^{i\mathbf{k} \cdot \boldsymbol{\vartheta}} \quad (n \in \mathbb{N}_0 = \{0, 1, 2, \dots\}); \quad (1.2)$$

the *triangular* one is

$$S_{nd}^{[t]}(g, \boldsymbol{\vartheta}) := \sum_{|\mathbf{k}|_1 \leq n} \hat{g}(\mathbf{k}) e^{i\mathbf{k} \cdot \boldsymbol{\vartheta}} \quad (n \in \mathbb{N}_0). \quad (1.3)$$

Above, $|\mathbf{k}|_\infty = \max_{1 \leq l \leq d} |k_l|$ and $|\mathbf{k}|_1 = \sum_{k=1}^d |k_l|$ (they are the l_p norms of the multiindex \mathbf{k}). The names “rectangular” and “triangular” refer to the shape of the corresponding indices of terms when $d = 2$ and $0 \leq k_1, k_2, |\mathbf{k}|_\infty \leq n, |\mathbf{k}|_1 \leq n$ respectively.

In a way the investigation of the $S_{nd}^{[r]}$ is apparent: in many cases in essence it is a one variable problem (see the above works [1,2]).

However there are only relatively few works dealing with the triangular (or l_1) summability: see Herriot [3].

In a recent paper [4] Berens and Xu prove the analogue of the famous Fejér paper on the $(C, 1)$ mean of $S_n(f)$ (here and later S_n stands for $S_{n1} \equiv S_{n1}^{[r]} \equiv S_{n1}^{[t]}$): actually it turns out that $(C, 2d - 1)$ in a way corresponds to $(C, 1)$ for any $d \geq 1$. Moreover the explicit formula for the corresponding Dirichlet kernel D_{nd} is crucial: it gives the possibility to prove many theorem for $S_{nd} \equiv S_{nd}^{[t]}$ analogous to statements on S_n (this formula which actually gives D_{nd} as a divided difference of univariable function was developed earlier by Xu [5]).

Exploiting this relation we prove some theorems on S_{nd} ; as it turns out they can be used to investigate $S_{nd}^{[r]}$, too.

2. New results

2.1

Introducing the notations

$$D_{nd}(\boldsymbol{\vartheta}) = \sum_{|\mathbf{k}|_1 \leq n} e^{i\mathbf{k} \cdot \boldsymbol{\vartheta}} \quad (n \geq 1), \quad (2.1)$$

where $\mathbf{k} \in \mathbb{Z}^d$, one can see that

$$\begin{aligned} S_{nd}(g, \boldsymbol{\vartheta}) &= (g * D_{nd})(\boldsymbol{\vartheta}) := \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} g(\boldsymbol{\vartheta} - \mathbf{t}) D_{nd}(\mathbf{t}) d\mathbf{t} \\ &= \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} g(\boldsymbol{\vartheta} + \mathbf{t}) D_{nd}(\mathbf{t}) d\mathbf{t}, \end{aligned} \quad (2.2)$$

where as before, $g \in C(\mathbb{T}^d)$, $\boldsymbol{\vartheta}, \mathbf{t} \in \mathbb{T}^d$ (cf. (1.3) or [2, Chs I, VII], [4]).

Let $\|g\| := \max_{\boldsymbol{\vartheta} \in \mathbb{T}^d} |g(\boldsymbol{\vartheta})|$,

$$\|S_{nd}\| := \max_{\substack{g \in C(\mathbb{T}^d) \\ \|g\| \leq 1}} \|S_{nd}(g, \boldsymbol{\vartheta})\| \quad (n \geq 1)$$

and

$$\|g\|_p := \left(\int_{\mathbb{T}^d} |g(\boldsymbol{\vartheta})|^p d\boldsymbol{\vartheta} \right)^{1/p}$$

if $g \in L^p := \{\text{the set of all measurable } 2\pi \text{ periodic (in each variable) functions on } \mathbb{T}^d\}$, $1 \leq p < \infty$.

We state

Theorem 2.1. *We have, for any fixed $d \geq 1$,*

$$\|D_{nd}\|_1 = \|S_{nd}\| \sim (\log n)^d \quad (n \geq 2).^1 \quad (2.3)$$

Remark. The general case of the upper estimation (when $d \geq 4$) is due to Professor Gábor Halász [6]. We shall give another argument applying a formula of Xu.

The lower estimation also belongs to Gábor Halász [6]. The proof is based on his original argument; we thank him for communicating the ideas.

2.2

One of the most characteristic properties of the Fourier series in one dimension is the so-called Faber–Marcinkiewicz–Berman theorem, namely that the operator S_n has the smallest norm among all projection operators (cf. [7, p. 281] for other details). This part extends the above statement for S_{nd} , $d \geq 1$.

Let \mathcal{T}_{nd} be the space of trigonometric polynomials of form

$$\sum_{|\mathbf{k}|_1 \leq n} (a_{\mathbf{k}} \cos(\mathbf{k} \cdot \boldsymbol{\vartheta}) + b_{\mathbf{k}} \sin(\mathbf{k} \cdot \boldsymbol{\vartheta})),$$

where $\mathbf{k} = (k_1, k_2, \dots, k_d)$ and $k_1, \dots, k_d \geq 0$, arbitrary real numbers. Moreover, let T_{nd} be a linear trigonometric projection operator on $C(\mathbb{T}^d)$, i.e. $T_{nd}(g, \boldsymbol{\vartheta}) = g(\boldsymbol{\vartheta})$ for $g \in \mathcal{T}_{nd}$ and $T_{nd}(g, \boldsymbol{\vartheta}) \in \mathcal{T}_{nd}$ for other $g \in C(\mathbb{T}^d)$.

Theorem 2.2. *For any linear trigonometric projection operator T_{nd} , one has*

$$\frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} T_{nd}(g_{\mathbf{t}}, \boldsymbol{\vartheta} - \mathbf{t}) d\mathbf{t} = S_{nd}(g, \boldsymbol{\vartheta}) \quad (g \in C(\mathbb{T}^d)), \quad (2.4)$$

$$\|T_{nd}\| \geq \|S_{nd}\|, \quad (2.5)$$

where $g_{\mathbf{t}}(\boldsymbol{\vartheta}) = g(\boldsymbol{\vartheta} + \mathbf{t})$ is the \mathbf{t} -translation operator.

¹ Here and later $a_n \sim b_n$ means that $0 < c_1 \leq a_n b_n^{-1} \leq c_2$ where c, c_1, c_2, \dots are positive constants, not depending on n ; they may denote different values in different formulae.

2.3

Here we formulate an analogue of [Theorem 2.2](#) which can be applied for the norm estimation of projection onto the space Π_{nd} (the algebraic analogue of \mathcal{T}_{nd} ; cf. [7, p. 284]). Let T_{nd} be a linear trigonometric projection operator defined on the even functions in $C(\mathbb{T}^d)$. We extend it to all of $C(\mathbb{T}^d)$ by defining $T_{nd}^*(g) := T_{nd}(G)$, where $2G(\boldsymbol{\vartheta}) = g(\boldsymbol{\vartheta}) + g(-\boldsymbol{\vartheta})$. Obviously, $\|T_{nd}^*\| = \|T_{nd}\|$.

Theorem 2.3. *We have for $g \in C(\mathbb{T}^d)$*

$$\frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} T_{nd}^*(g\mathbf{t} + g_{-\mathbf{t}}, \boldsymbol{\vartheta} - \mathbf{t}) d\mathbf{t} = S_{nd}^*(g, \boldsymbol{\vartheta}) = S_{nd}(G, \boldsymbol{\vartheta}), \quad (2.6)$$

$$\|T_{nd}\| = \|T_{nd}^*\| \geq \frac{1}{2} \|S_{nd}^*\| = \frac{1}{2} \|S_{nd}\|. \quad (2.7)$$

A simple consequence is

Theorem 2.4. *If P_{nd} is a projection of $C(I^d)$ onto Π_{nd} then*

$$\|P_{nd}\| \geq \frac{1}{2} \|S_{nd}\|. \quad (2.8)$$

Above, P_{nd} is a projection of $C(I^d)$ ($:=$ the set of continuous functions of d -variables on $I^d = [-1, 1]^d$) onto Π_{nd} iff it is linear, $P_{nd}(p) = p$ if $p \in \Pi_{nd}$ and $P_{nd}(f) \in \Pi_{nd}$ for any $f \in C(I^d)$.

As an application of [Theorem 2.4](#), see the papers [8–10] dealing with the two-dimensional Lagrange interpolation defined on special node systems.

Remark. Using our methods we intend to settle the “critical cases” in the paper [11, p. 290.] in our forthcoming work.

3. Proofs

3.1. Proof of [Theorem 2.1](#)

3.1.1

The relation

$$\|D_{nd}\|_1 = \|S_{nd}\| \quad (3.1)$$

is a simple consequence of the Riesz representation theorem (see [12, IV. 6.3] or [13]).

3.1.2

Next, we prove the relation

$$\|D_{nd}\|_1 \sim (\log n)^d \quad (n \geq 2).$$

Our basic tool is the relation proved by Xu (cf. [5, Lemma 1])

$$D_{nd}(\boldsymbol{\vartheta}) = (-1)^{\left[\frac{d-1}{2}\right]} \sum_{l=1}^d \frac{2 \cos \frac{\vartheta_l}{2} (\sin \vartheta_l)^{d-2} \text{soc} \frac{2n+1}{2} \vartheta_l}{\prod_{\substack{j=1 \\ j \neq l}}^d (\cos \vartheta_l - \cos \vartheta_j)}, \quad (3.2)$$

where the function soc (sin or cos) is defined by

$$\text{soc} \vartheta := \begin{cases} \sin \vartheta & \text{if } d \text{ is odd,} \\ \cos \vartheta & \text{if } d \text{ is even.} \end{cases} \quad (3.3)$$

3.1.3

First we prove that $\|D_{nd}\|_1 \leq c(\log n)^d$. Indeed, let

$$e_d := \left\{ \boldsymbol{\vartheta} : |\vartheta_k| \leq n^{-d}, |\vartheta_k - \vartheta_l| \leq n^{-d}, 1 \leq k \neq l \leq d \right\}. \quad (3.4)$$

Obviously $|e_d| \leq A(d)n^{-d}$, where $A(d) > 0$, fixed. If $E_d := \mathbb{T}^d \setminus e_d$ we get

$$\int_{\mathbb{T}^d} |D_{nd}| = \int_{e_d} |D_{nd}| + \int_{E_d} |D_{nd}|. \quad (3.5)$$

By (2.1), $|D_{nd}(\boldsymbol{\vartheta})| \leq 2^d \binom{2n+1}{d} \leq B(d)n^d$, whence

$$\int_{e_d} |D_{nd}| \leq A(d)B(d).$$

At the second integral of (3.5) we estimate only one term on the right-hand side of (3.2) on $F_d := E_d \cap [0, \frac{\pi}{2}]^d$ and then we use symmetry. By

$$\sin \vartheta_1 \leq 2 \sin \frac{\vartheta_1 + \vartheta_j}{2} \quad (2 \leq j \leq d, \boldsymbol{\vartheta} \in F_d) \quad (3.6)$$

(consider the cases $\vartheta_1 \leq \vartheta_j$ or $\vartheta_1 > \vartheta_j$) and

$$\cos \vartheta_1 - \cos \vartheta_j = 2 \sin \frac{\vartheta_1 + \vartheta_j}{2} \sin \frac{\vartheta_1 - \vartheta_j}{2}, \quad (3.7)$$

we get an estimate for s_1 (the first term of (3.2)) as follows:

$$\begin{aligned} |s_1| &\leq c \cdot \frac{\prod_{j=2}^d \sin \frac{\vartheta_1 + \vartheta_j}{2}}{\sin \vartheta_1 \prod_{j=2}^d \sin \frac{\vartheta_1 + \vartheta_j}{2} \left| \sin \frac{\vartheta_1 - \vartheta_j}{2} \right|} \\ &\leq \frac{c_1}{\sin \vartheta_1 \prod_{j=2}^d \left| \sin \frac{\vartheta_1 - \vartheta_j}{2} \right|} \quad (\boldsymbol{\vartheta} \in F_d). \end{aligned}$$

Then, using the substitution $u_1 = \vartheta_1$, $u_j = \vartheta_j - \vartheta_1$, whence $\vartheta_1 = u_1$, $\vartheta_j = u_1 + u_j$ ($2 \leq j \leq d$), we get

$$\begin{aligned} \int_{F_d} |s_1| d\boldsymbol{\vartheta} &\leq c_1 \int_{F_d} \frac{1}{\sin \vartheta_1 \prod_{j=2}^d \left| \sin \frac{\vartheta_1 - \vartheta_j}{2} \right|} d\boldsymbol{\vartheta} \\ &\leq c_2 V_d \int_{\frac{1}{n^d}}^{\frac{\pi}{2}} \int_{\frac{1}{n^d}}^{\frac{\pi}{2}} \cdots \int_{\frac{1}{n^d}}^{\frac{\pi}{2}} \frac{1}{\prod_{j=1}^d u_j} du_1 du_2 \dots du_d \\ &= c_2 \left(\log \frac{\pi n^d}{2} \right)^d \sim (\log n)^d, \end{aligned}$$

where if

$$U_d := \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 1 & 1 & 0 & \cdots & 0 \\ 1 & 0 & 1 & \cdots & 0 \\ & & \cdots & & \\ 1 & 0 & 0 & \cdots & 1 \end{pmatrix}$$

is the matrix of the transformation, then $V_d = |\det U_d| = 1$.

3.1.4

Here we prove that

$$\|D_{nd}\|_1 \geq c(\log n)^d \quad (n \in \mathbb{N}) \quad (3.8)$$

with a constant $c > 0$ independent of n .

Our proof is based on Fejér's classical example (see [14, Vol. 2, Ch. 2/1]). We suggest the interested reader to examine it carefully.

For every $n \in \mathbb{N}$ we shall construct a trigonometric polynomial $f_n(\mathbf{t}) = \sum_{\mathbf{j}} c_{\mathbf{j}} e^{i\mathbf{j} \cdot \mathbf{t}}$ with

$$\|f_n\| \leq 1 \quad \text{and} \quad |S_{nd}(f_n, \mathbf{0})| \geq c(\log n)^d \quad (n \in \mathbb{N}), \quad (3.9)$$

where $c > 0$ is independent of n .

With these polynomials we have

$$\|D_{nd}\|_1 = \|S_{nd}\| = \max_{\substack{g \in C(\mathbb{T}^d) \\ \|g\| \leq 1}} \|S_{nd}(g, \cdot)\| \geq |S_{nd}(f_n, \mathbf{0})|$$

which proves (3.8).

For the construction of f_n first we choose real numbers α_j, β_j ($j = 1, 2, \dots, d$) for which

$$1 > \alpha_1 > \beta_1 > \alpha_2 > \beta_2 > \cdots > \alpha_d > \beta_d > 0. \quad (3.10)$$

Starting from Fejér's classical example \mathcal{F}_m (see (3.12)) let us consider the trigonometric polynomials

$$F_j(t) := \sum_{|k_j|=n^{\beta_j}}^{[n^{\alpha_j}]} \frac{1}{k_j} e^{ik_j t} \quad (t \in [0, 2\pi), 1 \leq j \leq d). \quad (3.11)$$

As we know for the trigonometric polynomials

$$\mathcal{F}_m(t) := \sum_{0 < |l| \leq m} \frac{1}{l} e^{ilt} \quad (t \in \mathbb{R}, m \in \mathbb{N}) \quad (3.12)$$

we have

$$|\mathcal{F}_m(t)| = 2 \left| \sum_{l=1}^m \frac{\sin lt}{l} \right| \leq 4\sqrt{\pi} \quad (t \in \mathbb{R}, m \in \mathbb{N})$$

(see [15,16], [14, Vol. I, (118)]). Therefore we get

$$\begin{aligned} |F_j(t)| &= |\mathcal{F}_{[n^{\alpha_j}]}(t) - \mathcal{F}_{[n^{\beta_j]-1]}(t)| \leq |\mathcal{F}_{[n^{\alpha_j}]}(t)| + |\mathcal{F}_{[n^{\beta_j]-1]}(t)| \\ &\leq 8\sqrt{\pi} =: M. \end{aligned} \quad (3.13)$$

Using $\mathbf{e}_j := (0, \dots, 0, 1, 0, \dots, 0) (1 \leq j \leq d)$ (the canonical unit vectors of \mathbb{R}^d) we define the polynomial $g_n(\mathbf{t}) := M^d f_n(\mathbf{t})$ as follows

$$\begin{aligned} g_n(\mathbf{t}) &:= e^{i n \mathbf{e}_d \cdot \mathbf{t}} \cdot F_d(-\mathbf{e}_d \cdot \mathbf{t}) \cdot \prod_{j=1}^{d-1} F_j((\mathbf{e}_j - \mathbf{e}_d) \cdot \mathbf{t}) \\ &= e^{i n t_d} \sum_{|k_d|=n^{\beta_d}}^{[n^{\alpha_d}]} \frac{e^{-i k_d t_d}}{k_d} \prod_{j=1}^{d-1} \left(\sum_{|k_j|=n^{\beta_j}}^{[n^{\alpha_j}]} \frac{1}{k_j} e^{i(k_j t_j - k_j t_d)} \right), \end{aligned} \quad (3.14)$$

where $\prod_{j=1}^0 \cdots := 1$ and $(\mathbf{e}_j - \mathbf{e}_d) \cdot \mathbf{t} = t_j - t_d$.

Using (3.13) we obtain that $|f_n(\mathbf{t})| \leq 1 (t \in \mathbb{T}^d)$, i.e. the first requirement of (3.9) holds.

The polynomial $g_n(\mathbf{t})$ can be written as

$$g_n(\mathbf{t}) = \sum_{k_1, \dots, k_d}^* \frac{1}{k_1 k_2 \cdots k_d} e^{i \mathbf{k} \cdot \mathbf{t}},$$

where $\mathbf{k} := (k_1, k_2, \dots, k_{d-1}, n - k_1 - k_2 - \cdots - k_d)$ and the notation \sum_{k_1, \dots, k_d}^* means that we take the summation for indices $[n^{\beta_j}] \leq |k_j| \leq [n^{\alpha_j}] (1 \leq j \leq d)$.

The triangular partial sum of the Fourier series of g_n at the point $\mathbf{0}$ is given by

$$S_{nd}(g_n, \mathbf{0}) = \sum_{\substack{k_1, \dots, k_d \\ |\mathbf{k}|_1 \leq n}}^* \frac{1}{k_1 k_2 \cdots k_d}. \quad (3.15)$$

Now we prove that in this sum appear *only the positive indices* k_1, \dots, k_d .

First we remark that $n - k_1 - k_2 - \dots - k_d \geq 0$. Then, by $|k_j| - k_j \geq 0$ ($1 \leq j \leq d$), we get

$$\begin{aligned} n &\geq |\mathbf{k}|_1 = |k_1| + |k_2| + \dots + |k_{d-1}| + |n - k_1 - k_2 - \dots - k_d| \\ &= \sum_{j=1}^{d-1} (|k_j| - k_j) + n - k_d \geq n - k_d, \end{aligned}$$

whence we obtain that $k_d \geq 0$. Applying (3.10) we have $k_d > 0$.

Let us suppose that for a fixed index j^* ($1 \leq j^* \leq d-1$) we have $k_{j^*} < 0$. Since $-k_{j^*} - k_d > 0$ for $n \geq n_0$ (see (3.10)), we would get

$$|\mathbf{k}|_1 = \sum_{j=1}^{d-1} (|k_j| - k_j) + n - k_d \geq -2k_{j^*} + n - k_d > n - k_{j^*} > n$$

— a contradiction. This means that in (3.15) appear only positive indices k_1, k_2, \dots, k_{d-1} , indeed. Therefore we get (see (3.14))

$$S_{nd}(f_n, \mathbf{0}) = \frac{1}{M^d} \prod_{j=1}^d \left(\sum_{k_j=[n^{\beta_j}]}^{[n^{\alpha_j}]} \frac{1}{k_j} \right) \geq c(\log n)^d$$

which proves the second requirement of (3.9).

Thus the proof of (3.8) is complete. \square

3.2. Proof of Theorem 2.2.

It is analogous to the classical argument (cf. [7, p. 282]).

3.2.1

First we verify that the integrand in (2.4) is a continuous function of $(\boldsymbol{\vartheta}, \mathbf{t}) \in \mathbb{T}^d \times \mathbb{T}^d$. Indeed, if $\boldsymbol{\vartheta}_0, \mathbf{t}_0$ are fixed then

$$\begin{aligned} |T_{nd}(g_{\mathbf{t}}, \boldsymbol{\vartheta}) - T_{nd}(g_{\mathbf{t}_0}, \boldsymbol{\vartheta}_0)| &\leq |T_{nd}(g_{\mathbf{t}}, \boldsymbol{\vartheta}) - T_{nd}(g_{\mathbf{t}_0}, \boldsymbol{\vartheta})| \\ &\quad + |T_{nd}(g_{\mathbf{t}_0}, \boldsymbol{\vartheta}) - T_{nd}(g_{\mathbf{t}_0}, \boldsymbol{\vartheta}_0)| =: \delta_1 + \delta_2. \end{aligned}$$

Above, $\delta_1 \leq \|T_{nd}\| \|g_{\mathbf{t}} - g_{\mathbf{t}_0}\| < \varepsilon$ if $\|\mathbf{t} - \mathbf{t}_0\|$ is small; on the other hand if $\|\boldsymbol{\vartheta} - \boldsymbol{\vartheta}_0\|$ is small then $\delta_2 < \varepsilon$, too.

So the integral (2.4) is well defined. Denoting it by $A(g, \boldsymbol{\vartheta})$ we prove that $A(g, \boldsymbol{\vartheta}) = S_{nd}(g, \boldsymbol{\vartheta})$ for any $g \in C(\mathbb{T}^d)$.

First, we verify that $A(g)$ is linear trigonometric projection operator. It is enough to prove the boundedness of $A(g)$. But it follows by

$$\begin{aligned} |A(g, \boldsymbol{\vartheta})| &\leq \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} |T_{nd}(g_{\mathbf{t}}, \boldsymbol{\vartheta} - \mathbf{t})| d\mathbf{t} \\ &\leq \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} \|T_{nd}\| \|g_{\mathbf{t}}\| d\mathbf{t} = \|T_{nd}\| \|g\|, \end{aligned}$$

whence $\|A\| \leq \|T_{nd}\|$.

3.2.2

Moreover, we verify that

(a) for $p \in \mathcal{T}_{nd}$, $A(p) = S_{nd}(p)$. Indeed, by $p_{\mathbf{t}}(\boldsymbol{\vartheta}) \in \mathcal{T}_{nd}$ we have that $T_{nd}(p_{\mathbf{t}}, \boldsymbol{\vartheta} - \mathbf{t}) = p_{\mathbf{t}}(\boldsymbol{\vartheta} - \mathbf{t}) = p(\boldsymbol{\vartheta})$, whence by (2.4), $A(p) = p$, which by $S_{nd}(p) = p$ (cf. [7, p.209]), gives $A(p) = S_{nd}(p)$ if $p \in \mathcal{T}_{nd}$.

(b) Now let $p(\boldsymbol{\vartheta}) = \cos(\mathbf{m} \cdot \boldsymbol{\vartheta})$, $|\mathbf{m}|_1 > n$. We prove that $A(p, \boldsymbol{\vartheta}) = \mathbf{0}$. Indeed, by

$$p_{\mathbf{t}}(\boldsymbol{\vartheta}) = \cos(\mathbf{m} \cdot \boldsymbol{\vartheta}) \cos(\mathbf{m} \cdot \mathbf{t}) - \sin(\mathbf{m} \cdot \boldsymbol{\vartheta}) \sin(\mathbf{m} \cdot \mathbf{t})$$

(to verify, see the exponential form, say) we get

$$\begin{aligned} (2\pi)^d A(p, \boldsymbol{\vartheta}) &= \int_{\mathbb{T}^d} T_{nd}(p, \boldsymbol{\vartheta} - \mathbf{t}) \cos(\mathbf{m} \cdot \mathbf{t}) d\mathbf{t} \\ &\quad - \int_{\mathbb{T}^d} T_{nd}(q, \boldsymbol{\vartheta} - \mathbf{t}) \sin(\mathbf{m} \cdot \mathbf{t}) d\mathbf{t} =: I_1 + I_2, \end{aligned}$$

where $q(\boldsymbol{\vartheta}) = \sin(\mathbf{m} \cdot \boldsymbol{\vartheta})$. Consider the first integral, say. By definition, $T_{nd}(p, \boldsymbol{\vartheta} - \mathbf{t})$ is a linear combination of terms having the form $\cos(\mathbf{k} \cdot \mathbf{t})$ ($|\mathbf{k}|_1 \leq n$), whence in I_1 we have integrals of type

$$\int_{\mathbb{T}^d} \cos(\mathbf{k} \cdot \mathbf{t}) \cos(\mathbf{m} \cdot \mathbf{t}) d\mathbf{t}. \quad (3.16)$$

But they are equal to zero. Indeed, as it is well known (and easy to see) that the functions $\{e^{i\mathbf{r} \cdot \mathbf{t}}; \mathbf{r} \in \mathbb{Z}_0^d, \mathbf{t} \in \mathbb{T}^d\}$ are mutually orthogonal, i.e.

$$\int_{\mathbb{T}^d} e^{i\mathbf{r} \cdot \mathbf{t}} e^{-i\mathbf{s} \cdot \mathbf{t}} d\mathbf{t} \neq 0 \quad \text{iff } \mathbf{r} = \mathbf{s};$$

whence by

$$\begin{aligned} 2 \cos(\mathbf{r} \cdot \boldsymbol{\vartheta}) &= e^{i\mathbf{r} \cdot \boldsymbol{\vartheta}} + e^{-i\mathbf{r} \cdot \boldsymbol{\vartheta}}, \\ 2i \sin(\mathbf{r} \cdot \boldsymbol{\vartheta}) &= e^{i\mathbf{r} \cdot \boldsymbol{\vartheta}} - e^{-i\mathbf{r} \cdot \boldsymbol{\vartheta}} \end{aligned}$$

(one may verify them by induction for d), the functions

$$\{\mathbf{e}, \cos(\mathbf{r} \cdot \mathbf{t}), \sin(\mathbf{s} \cdot \mathbf{t})\};$$

$\mathbf{r}, \mathbf{s} \in \mathbb{N}^d$, $\mathbf{e} = \{1\}^d$ are mutually orthogonal, too. That means, in (3.16) using $|\mathbf{k}|_1 < |\mathbf{m}|_1$ we get $\mathbf{k} \neq \mathbf{m}$, i.e. the integral is zero, indeed.

The other cases are similar. Using that $S_{nd}(p) = S_{nd}(q) = 0$ ($|\mathbf{m}|_1 > n$), we get $A(p) = A(q) = S_{nd}(p) = S_{nd}(q) = 0$ at case (b).

3.2.3

Summarizing, we obtained that $S_{nd}(g) = A(g)$ for all trigonometric polynomials, whatever their degree. Using that they form a dense set in $C(\mathbb{T}^d)$ (cf. [2, Theorem 1.7, p. 248]), the operators A and S_{nd} coincide. \square

3.2.4

Now (2.5) is immediate: By (2.4) we write

$$|S_{nd}(g, \boldsymbol{\vartheta})| \leq \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} |T_{nd}(g_{\mathbf{t}}, \boldsymbol{\vartheta} - \mathbf{t})| d\mathbf{t} \leq \|T_{nd}\| \|g\|. \quad \square$$

3.3. Proof of Theorem 2.3.

It is very similar to the previous one. For the sake of variety we use the system $\{e^{i\mathbf{r}\cdot\mathbf{t}}\}$.

Let $g(\boldsymbol{\vartheta}) = e^{i\mathbf{k}\cdot\boldsymbol{\vartheta}}$, $|\mathbf{k}|_1 \leq n$. Then by $g_{\pm\mathbf{t}}(\boldsymbol{\vartheta}) = e^{i\mathbf{k}\cdot(\boldsymbol{\vartheta}\pm\mathbf{t})}$, whence

$$T_{nd}^*(g_{\mathbf{t}} + g_{-\mathbf{t}}, \boldsymbol{\vartheta} - \mathbf{t}) = 2 \cos(\mathbf{k} \cdot (\boldsymbol{\vartheta} - \mathbf{t})) \cos(\mathbf{k} \cdot \mathbf{t}).$$

So the integrand in (2.6) is equal to $\cos(\mathbf{k} \cdot \boldsymbol{\vartheta})$ (simple calculation) which is also the right-hand side of (2.6).

(b) If $g(\boldsymbol{\vartheta}) = e^{i\mathbf{m}\cdot\boldsymbol{\vartheta}}$, $|\mathbf{m}|_1 > n$, then the integrand is $T(\boldsymbol{\vartheta} - \mathbf{t}) \cos(\mathbf{m} \cdot \mathbf{t})$, where $T \in \mathcal{T}_{nd}$, i.e. the left-hand side is zero, which holds for the right-hand side, too.

Using the above facts, we can get (2.6) as before (cf. 3.2.1–3.2.3).

Finally, using the obvious fact $\|T_{nd}\| = \|T_{nd}^*\|$, (2.7) easily comes from (2.6) (see 3.2.4). \square

3.4. Proof of Theorem 2.4.

The map

$$V : f(\mathbf{x}) \rightarrow f(\cos \boldsymbol{\vartheta}) \equiv g(\boldsymbol{\vartheta})$$

is an isometric operator with $\|V\| = 1$. If we define the projection in Theorem 2.3 by

$$T_{nd} = V P_{nd} V^{-1},$$

then by (2.7)

$$\frac{1}{2} \|S_{nd}\| \leq \|T_{nd}^*\| = \|T_{nd}\| = \|V P_{nd} V^{-1}\|,$$

as it was stated. \square

Acknowledgments

The authors thank the referees for their comments and suggestions. At the same time we express again our sincere thanks to Professor Gábor Halász. The work was supported by the Hungarian National Scientific Research Foundation (OTKA) under Grant Ns. T37299 and T047132.

References

- [1] A. Zygmund, Trigonometric Series, vols. I and II, Cambridge University Press, Cambridge, 1959.
- [2] E.M. Stein, G. Weiss, Introduction to Fourier Analysis on Euclidean Spaces, Princeton University Press, Princeton, 1971.
- [3] J.G. Herriot, Nörlund summability of multiple Fourier series, Duke Math. J. 11 (1944) 735–754.
- [4] H. Berens, Y. Xu, Fejér means for multivariate Fourier series, Mat. Z. 221 (1996) 449–465.
- [5] Y. Xu, Christoffel functions and Fourier series for multivariate orthogonal polynomials, J. Approx. Theory 82 (2) (1995) 205–239.
- [6] G. Halász, personal communications.
- [7] R.A. DeVore, G.G. Lorentz, Constructive Approximation, Springer-Verlag, Berlin, 1993.
- [8] L. Bos, S. De Marchi, M. Vianello, On Lebesgue constant of Xu interpolation formula, J. Approx. Theory 141 (2006) 134–141.
- [9] B. Della Vecchia, G. Mastroianni, P. Vértési, Exact order of the Lebesgue constants for bivariate Lagrange interpolation at certain node-systems, Studia Sci. Math. Hungar. (in press).
- [10] Y. Xu, Lagrange interpolation on Chebyshev points of two variables, J. Approx. Theory 87 (2) (1996) 220–238.

- [11] Z. Li, Y. Xu, Summability of product Jacobi expansions, *J. Approx. Theory* 104 (2000) 287–301.
- [12] N. Dunford, J.T. Schwartz, *Linear Operators, Part I: General Theory*, John Wiley & Sons, New York, 1988.
- [13] K. Yosida, *Functional Analysis*, Springer–Verlag, 1965.
- [14] I.P. Natanson, *Constructive Function Theory*, vol. 1–3, Ungar, New York, 1964–1965.
- [15] L. Fejér, Sur les singularités de la série de Fourier des fonctions continues. (French), *Ann. Sci. École Norm. Sup.* 28 (1911) 63–104.
- [16] L. Fejér, A függvény szakadásának meghatározása Fourier-féle sorából (Hungarian), *Mat. és Term. Értesítő* 31 (1913) 385–415.